



Available online at <https://www.s-epub.in/>

Pharma Research

An International Peer Reviewed, Indexed Journal

10.62655/s-epub.2021.v13.i02.pp1-7

Stability Indicating Analytical Method Development, Validation, Method Transfer And Impurity Profile (Related Substances) of 2,4-Dihydroxy-5-Fluoropyrimidine by Liquid Chromatography

S.Imam Pasha^{1*},

Affiliation

^{*1}Department Of Pharmaceutical Analysis & Quality Assurance, Sultan-UL-Uloom College Of Pharmacy, Banjara Hills, Hyderabad-500034, TS, India

Article info

Published on: 21-10-2021

ISSN : 0975-8216

Abstract:

Uracil, 5-Methoxy Uracil, 5-Chloropyrimidine-2,4(1H,3H)-dione, Dihydropyrimidine-2,4,5(3H)-trione, and other related chemicals were the subject of a stability-indicating liquid chromatographic method's development. Using a column with dimensions of 250mm × 5µm (YMC Pack ODS AQ), a flow rate of 1.0 mL/min at a wavelength of 266 nm, an injection volume of 20 µL, and a run period of 30 minutes, validation was carried out on a greater concentration of 2,4-dihydroxy-5-fluoro pyrimidine.

I INTRODUCTION

The anti-metabolite 2, 4-dihydroxy-5-fluoropyrimidine is the drug of choice for cancer chemotherapy because of its irreversible effect on the enzyme thymidylate synthase.[2] is cited. Cells are targeted at particular stages of the cell cycle. 1, 3, 7, and the contaminants such The following impurities were commonly found in 2,4-dihydroxy-5-fluoropyrimidine in bulk drug and formulation sites: pyrimidine-2,4,6 (1H,3H,5H)-trione,

dihydropyrimidine-2,4,5(3H)-trione, unacil, 5-chloropyrimidine-2,4(1H,3H)-dione. No HPLC method was found in the literature to identify or assess the limit of these five impurities. The approach was subjected to forced degradation investigations, which confirmed its stability. It has great potential for use in bulk drug facilities and formulation sites, where the limit of contaminants may be determined.

II MATERIALS & METHODS

Details of Chemicals: Fluorouracil Standard (Batch IOG371, USP Grade), Fluorouracil 50mg/ml (Batch IFU-319(B), Ingénues), Mono

basic Potassium Phosphate (Batch QF4Q641420, Merck), Acetonitrile (Batch :IA51F65025, Merck)

Instruments (Columns, serial no.):

HPLC: VLS-DR/HPLC/05

VLS-DR/HPLC/12

VLS-DR/HPLC/16

VLS-DR/HPLC/17

Analytical Balance: VLS-DR/BAL/01

pH Meter: VLS-DR/PHM/01

Description of Analytical Method:

Chromatographic Parameters:

Column: 250mm, 5µm (YMC Pack ODS AQ or Phenomenex Luna C18 (2) 100A or equivalent to L1)

Flow rate: 1.0 mL/min

Wavelength: 266 nm

Injection Volume: 20 µL

Column oven Temperature: Ambient

Run time : 30 minutes

Preparation of Mobile Phase

Weigh about 6.8 g of Monobasic Potassium phosphate and transfer into 1000 ml of water, dissolve and adjust the pH of this solution to 5.7 with 5M Potassium hydroxide.

Fluorouracil Standard Stock solution

Accurately weigh and transfer 5 mg of Fluorouracil standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluent. Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluent.

Fluorouracil Related Compound A Stock solution (Stock-A)

Weigh and transfer 5 mg of Fluorouracil Impurity-A standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluent. Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluent.

Fluorouracil Related Compound B Stock solution (Stock-B)

Weigh and transfer 5 mg of Fluorouracil Impurity-B standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluent.

Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluent.

Fluorouracil Related Compound C Stock solution (Stock-C)

Weigh and transfer 5 mg of Uracil (Impurity-C) standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluents. Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluents.

Fluorouracil Related Compound D Stock solution (Stock-D)

Weigh and transfer 5 mg of Fluorouracil Impurity-D standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluents. Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluent.

Fluorouracil Related Compound E Stock solution (Stock-E)

Weigh and transfer 5 mg of Fluorouracil Impurity-E standard into a 25 mL volumetric flask, dissolve and dilute to volume with diluent. Transfer 1 mL of this solution into a 100 mL volumetric flask and dilute to volume with diluent.

Preparation of Standard solution

Transfer each 1.0 mL of Fluorouracil Standard Stock solution, Stock A, B, C, D, E into a 10 mL volumetric flask and dilute to volume with diluent and mix well.

Preparation of Test solution:

Transfer 1 mL of Fluorouracil injection (50mg/mL) into a 50 mL volumetric flask, dissolve and dilute to the volume with diluent. Transfer 1.0 mL of the above solution into a 10 mL volumetric flask and dilute to volume with diluent and mixed well.

System Suitability

Injected blank and standard solution for six times into the HPLC system. Specificity

Interference Study:

As per methodology, injected blank, placebo solution once each and standard Solution, sample solution and spiked solution and checked the peak interference of blank, placebo and impurities at the retention time of Fluorouracil and its Impurities. Prepared and injected each impurity at 1% level individually and checked the interference at each impurity retention time.

Precision:

System Precision

As per methodology, injected blank and standard solution six times and check standard once into HPLC system.

Method Precision

Analyzed six test preparations of Fluorouracil injection 50 mg/mL as per the methodology and determined the % RSD of any individual impurity and total impurities from six sample preparations of Fluorouracil.

Intermediate Precision: Determined
the Intermediate precision by preparing six test preparations of Fluorouracil injection 50 mg/mL as per the methodology and analyzed as per the test method by different analyst on different analyst on different day by using different system with different column. Here intermediate precision study was carried out at the receiving site. Intermediate precision which was performed as a co-validation (inter laboratory variation) and considered for method transfer activity.

Limit of Detection and limit of quantification

As per methodology, injected blank, reference solution for six times and then injected LOD & LOQ Solutions into HPLC.

Linearity

Linearity for Fluorouracil was determined in the concentration range from 50 to 150 % levels of test concentration levels.

Accuracy

As per methodology, prepared 50%, 100% and 150% sample solutions of Fluorouracil working concentration and demonstrated the accuracy on sample into HPLC. Calculated the system suitability parameters and % mean recovery.

Range

From the results of Method Precision, Linearity and Accuracy it was concluded that the range of the Analytical method was established from 50 to 150 % of target concentration.

Robustness:

Effect of Variation in Flow rate

System suitability preparations were analyzed as per the methodology at low column flow (0.9 mL/min) and high column flow (1.1 mL/min) variation in flow rate.

Acceptance criteria

The resolution between Fluorouracil and Uracil (Impurity-C) should be not less than 2.0 in the standard solution. The % RSD of the area of Fluorouracil peak from six replicate injections of standard solution should be not more than 5.0.

Effect of Variation in pH

System suitability preparations were analyzed as per the methodology at low pH (5.6) and high pH (5.8) variation in buffer.

Effect of Variation in Column Oven Temperature

System suitability preparations were analyzed as per the methodology at high column Oven temperature (30°C) variation in column Oven temperature.

III RESULTS & DISCUSSION

Table.1: Relative Retention Time of Impurities

Name	Relative Retention Time
Fluorouracil Related compound A, Pyrimidine-2,4,6(1H,3H,5H)-trione.	0.5
5-Fluorouracil Related compound B, Dihydropyrimidine-2,4,5(3H)-trione	0.7
Uracil	0.9
Fluorouracil	1.0
5-Methoxy Uracil	1.6
Fluorouracil Related compound E, 5-Chloropyrimidine-2,4(1H,3H)-dione	1.9

Table.2: Placebo Interference Data

S.No.	RT of Fluorouracil impurity A	RT of Fluorouracil impurity B	RT of Fluorouracil impurity C	RT of Fluorouracil impurity D	RT of Fluorouracil impurity E
Interference found (Yes/No)					
1	No	No	No	No	No

Table.3: Impurities Interference Data

S.No	Name	Interference Due to other Impurities(Yes/No)
1	Fluorouracil impurity A	No
2	Fluorouracil impurity B	No
3	Fluorouracil impurity C	No
4	Fluorouracil impurity D	No
5	Fluorouracil impurity E	No

Table.4: Interference from Degradation process in blank

Name Condition	Stress Condition	Interference at RT of Fluorouracil (Yes/No)
Acid	1.0 mL of 5 M HCl for 180 min at 60°C	No
Base	1.0 mL of 5 M NaOH for 180 min at 60°C	No
Peroxide	1.0 mL of 30 % H ₂ O ₂ for 5 min at 60°C	No
Water	1.0 mL of Water for 60 min at 90°C	No
Thermal	105°C for 6 hours	No
Humidity	90 % RH for 5 days	No
Photo Stability	1.2 million lux hours for white light and /200Watts for square meter for UV light	No

Table 5: Complete Degradation Data

S.No	Type of Stress	Assay (%w/w)	Purity 1Angle	Purity 1 Threshold	Peak Purity (Pass/Fail)
1	Acid	95.2	0.59	1.645	Pass
2	Base	93.9	0.64	1.309	Pass
3	Peroxide	97.9	1.02	2.456	Pass
4	Thermal	99.0	0.45	1.465	Pass
5	Humidity	96.4	0.39	1.239	Pass
6	Photo stability	98.9	0.73	1.875	Pass

Table 6: Method precision Results

Sample	Any Individual impurity (%w/w)	Total impurities (%w/w)
01	0.0096	0.0249
02	0.0097	0.0260
03	0.0098	0.0265
04	0.0093	0.0264
05	0.0099	0.0252
06	0.0093	0.0243
Average	0.0096	0.0256
S.D	0.0003	0.0009
%RSD	2.6	3.5

Table.Tab

Table.7: Limit of Detection and Limit of Quantification

Name	LOD (ppm)	LOQ (ppm)
Fluorouracil	0.0006	0.0014

Table.8: Precision at LOQ

Preparation	Area
1	907
2	904
3	899
4	944
5	870
6	921
Average	908
STDEV	24.5173
% RSD	2.7

Table.9: Accuracy at LOQ Level of Fluorouracil

Sample No.	Fluorouracil		
	Added	Found	% Recovery
1	0.00141	0.00136	96.45
2	0.00141	0.00138	97.87
3	0.00141	0.00125	88.65
Mean			94.3
Std.dev			4.964
% RSD			5.3

Table.10: Linearity Results of Fluorouracil

Level (%w/w)	Fluorouracil Concentration	Fluorouracil Peak Area
LOQ	0.0015	978
50	0.0101	5761
80	0.0161	9420
100	0.0201	11919
120	0.0241	14313
150	0.0302	17956
Correlation Coefficient	0.9998	

Table11: Accuracy of Fluorouracil (Assay)





Sample No	Spike level	Added (mg/mL)	Found (mg/mL)	'%' Recovery	'%' Mean recovery	%RSD
1	50%	0.01005	0.01002	99.66	99.8	0.1
2	50%	0.01005	0.01004	99.87		
3	50%	0.01005	0.01004	99.84		
1	100%	0.02010	0.02014	100.19	100.1	0.2
2	100%	0.02010	0.02007	99.82		
3	100%	0.02010	0.02016	100.26		
1	150%	0.03015	0.03032	100.56	100.3	0.3
2	150%	0.03015	0.03025	100.33		
3	150%	0.03015	0.03017	100.04		


Table12: Assay Sample solution stability results (RT and 2-8°C)

Parameter		Any individual impurity	% Difference from Initial	Total impurities	% Difference from Initial
Initial		0.0096	-	0.0249	-
Day-1	Sample at 2-8°C	0.0094	0.0002	0.0264	0.0015
	Sample at RT	0.0072	0.0024	0.0253	0.0004
Day-2	Sample at 2-8°C	0.0123	0.003	0.0323	0.0074
	Sample at RT	0.0131	0.0035	0.0280	0.0031


Table13: Robustness: Effect of Variation in Flow rate

Parameter	Resolution	% RSD
Low flow	4.6	1.1
High flow	5.1	0.3
Acceptance Criteria	NLT 2.0	NMT 5.0

Spectra.1: Typical chromatogram of Standard**Spectra.2:** Typical chromatogram of H₂O₂ stress Sample**Spectra.3:** Typical chromatogram of Thermal stress Sample**Spectra.4:** Typical chromatogram of Humidity stress Sample**Spectra.5:** Typical chromatogram of Photolytic stress Sample



GRAPH.1: Fluorouracil Linearity



IV CONCLUSION

Forced degradation studies of developed method for identifying the related substances of 2, 4 - dihydroxy -5-fluoropyrimidine were established. The present analytical method was validated for all the validation parameters and the developed analytical method meets the required acceptance criteria. the present analytical method proved to be stability indicating because the results were within the acceptance criteria both at transferring site and receiving site therefore the method transfer stands successful and can be used for regular analysis in pharmaceutical analysis & quality control departments for its intended purpose.

REFERENCES

- [1]. Review of approaches for developing validated stability-indicating assays (Bakhshi and Singh, 2003). Article published in the Journal of Pharmaceutical and Biomedical Research in 2002, volume 28, pages 1011-1040.

Stability testing of novel pharmaceutical compounds and products is covered in the ICH report [2]. The International Conference on Harmonisation (IFPMA) was held in Geneva in 2003. Photo stability assessment of novel medicinal compounds and products is described in [3]. ICH, Stability testing."International Conference on Harmonisation, IFPMA" (1996) occurred in Geneva.

This information is sourced from a 2004 publication by Gerber et al. [4]. "Practical aspects of fast reversed-phase high- performance liquid chromatography using 3µm particle packed columns and monolithic columns"

- [5].Effre, C. (2001). "Milestones in Chromatography: The Birth of Partition [6]. Chromatography", LCGC. 19 (5): 506-512 [online]. Accessed on 2016-02-02.
- [7].Author: Richard A. Henry; Date: February 1, 2009 "The Early Days of HPLC at DuPont" . Internet Resource for Chromatography. Avanstar Networks, Part II [8].Dynamics of Chromatography, Part I, J. Calvin Giddings (1965). Fundamentals and Concepts. (New York: Marcel Dekker, Inc., p. 281) [9].An RP-HPLC Assay Method for 5-Fluorouracil That Indicates Stability, by V. R. Sinha, *R. V. Kumar, and J. R. Bhinge, published in the Indian Journal of Pharmaceutical Sciences in 2009 (71(6): 630-637), was recently developed.