



Available online at <https://www.s-epub.in/>

Pharma Research

An International Peer Reviewed, Indexed Journal

10.62655/s-epub.2021.v13.i01.pp1-4

Impact of the aqueous extract of the bark of *Sclerocarya birrea* on the lipid membrane structure hepatocytes Wistar rats

Michodjehoun M. Clementine, Hoteyi Ismael,

Affiliation

Laboratory of Biomembrans and Signalling Cell University of Abomey-Calavi, Department of Animal Physiology, Rue Circulaire des Sciences, Campus d'Abomey-Calavi, Benin

Article info

Published on: 07-03-2021

ISSN: 0975-8216

KEYWORDS:

Sclerocarya birrea, bark, membran lipid, Benin

ABSTRACT:

The researchers in this work set out to analyse how different water-based extracts from the stem bark of *Sclerocarya birrea* affected the lipid membrane structure of hepatocytes in Wistar rats. We followed Bintou M.'s (2010) instructions to make an aqueous extraction using powdered dry bark of *Sclerocarya birrea* for this purpose. The biochemical testing of serum transaminase was the next step.

An increase in serum transaminase levels indicative of hepatocyte necrosis is not induced by aqueous extracts of *Sclerocarya birrea* bark, according to our findings. The analysis of rat hepatocyte membrane lipid profiles using thin layer chromatography of lipids verified this, showing that hepatocyte damage might occur at high dosages (≥ 200 mg/kg) of an aqueous extract of *Sclerocarya birrea* stem bark. Because there was no discernible difference in lipid profile between the negative control and doses of 50 and 100 mg/kg of *Sclerocarya birrea* aqueous extract, it is safe to assume that this extract is hazardous at higher concentrations. *Sclerocarya birrea* aqueous extract is hazardous at concentrations of 200 mg/kg or above.

INTRODUCTION

Plants in Africa are well-known for their medicinal properties and the diversity and abundance of the continent's flora. Among the 300,000 plant species in the world, more than 200,000 are native to

tropical African nations and have shown medicinal use (Sofowora 1993). Medicinal plants are an integral part of African health care. The vast majority of low-income populations in developing

nations rely on medicinal plants for both basic healthcare and food security; this makes them a priceless cultural artefact for all of mankind (Salhiet al., 2010). Medications are out of reach for many citizens in these nations due to their low incomes (Diallo, 2014). Conventional medicines, including herbal remedies, have a number of issues, the most significant of which is the lack of information on their efficacy, mechanism of action, active ingredients, doses, indications, safety, and quality control. According to Campa (2005), tropical plants provide over 80% of the natural chemicals utilised in biomedicine. The African governments must rethink their health policies, which have relied too heavily on pharmaceutical imports for far too long, and instead include traditional healing methods into their programmes. Serious chemical and biological decoding studies should be conducted on medicinal plants with a strong human identification and excellent therapeutic reputation in order to prove their safety and find more effective ways to utilise them. There have been several investigations on the benefits of the African native medicinal herb *Sclerocarya birrea*. Also, research has demonstrated that *Sclerocarya birrea* has antidiabetic properties (Coulibaly, 1988; Haidara, 1999; Fomba, 2001), that the plant's stem bark extract has glucose-lowering activity in rats (Ojewole 2003 ; Ojewole, 2004), that the plant's stem bark extracts have anti-diarrheal activities (Galves et al., 1991), and that the ester (-) - epicatechin isolated -3-galloyl from the plant's trunk bark has secretagogue activity (Galvez et al., 1992). These *Sclerocarya birrea* potentials pique our curiosity in investigating if there is a connection between the plant's active components and liver function, as the liver is a

A person's general well-being is impacted by the functioning of their regulatory body, which is also the primary organ responsible for detoxifying the body. The goal of this effort is therefore justified: "Impact of the aqueous extract of the bark of *Sclerocarya birrea* on the lipid membrane structure of the hepatocytes of Wistar rats."

I. MATERIALS AND METHODS

Plant Material

This is *Sclerocarya* trunk bark of *birrea*, harvested in July 2015 at the village of Gomez-kparou in the town of N'dali northeast Benin by a technical team in expédition. Les bark after receipt were washed and subjected to an oven at 40 ° C for 72h for drying. These dried peel were finely crushed and the powder obtained is then stored in glass jars to prevent the installation of

microorganisms pollutants.

Animal material

These liver Wistar type rats subjected to a sub-chronic gavage (14 days) and high at the pet of Biomembranes Laboratory and Cell Signaling FAST / UAC. Breeding is done in a lighted room 12 hours a day, and the temperature is ambient. The animals had free access to food and water.

Methods

Préparation of the aqueous extract of the bark of

To obtain the aqueous extract of the bark of the trunk *Sclerocarya birrea*, leaf powder 100g were weighed with an analytical balance Sartorius® and added to 1L of water, is brought to the boil for 30 minutes. The decoction is cooled for 20 minutes, filtered through gauze and cotton on. The resulting filtrate was overthrown in dishes and then put in the oven for evaporation at 50 ° C. After complete drying, the solids joined in the bottom plates are scraped with a spatula in stainless steel, then kept in bottles already labeled glass. The powders obtained were stored in glass vials, sterile and hermetically sealed.

For each series of experiments the extract is weighed and dissolved in distilled water to give solutions in the desired concentrations.

Gavage rats

In this handling, it is to administer the aqueous extracts of the barks of *Sclerocarya birrea* to Wistar rats (female) of about 200g of the same diet, and orally at various doses for about two weeks. These animals received no other medication in the time outside the extract.

- The rats were randomly divided into 06 lots, 03 rats (# 1, # 2 and # 3), then the doses per kg body weight per day are attributed to lots respectively, 0 mg / Kg; 50 mg / kg; 100 mg / kg; 200 mg / kg; 300 mg / kg; 400 mg / kg; and the volume is fixed to be administered at V = 01 ml.

- Then we weigh each rat of each batch to find the average weight and calculate the effective dose of extract to be administered to him. Then gave daily and at the same time the rats for 14 days.

- Lot 1 is the control group and receives oral administration of distilled water instead of the aqueous extract (0 mg / Kg) of *Sclerocarya birrea* for 14 days;

- Other lots 2; 3; 4; 5 and 6 are treated by oral administration at doses of 50 mg / kg; 100 mg / kg; 200 mg / kg; 300 mg / kg; 400 mg / kg body weight per day of the aqueous *Sclerocarya birrea* extract for 14 days

Figure 15: Picture of feeding a rat (Photo Amos Oluwagbemiga S., 2015).



- Finally, the rats were sacrificed 24 hours after the last feeding, to take our livers for various manipulations.

Isolation of rat hepatocytes by cold trypsinization technique [25; 26; 27; 28; 29]

The trypsinization is the method for disaggregating the tissue to isolate cells using the hot-exposure method or cold. Due to tryptic activity of trypsin in the hot method of trypsinization at 37 ° C; the tissue is damaged and harvesting the cells after 30 minutes incubation may lead to a reaggregation or damage cells. These inconveniences can be overcome by employing a minimum concentration of trypsin and a long incubation time to increase the efficiency of separation of trypsin. In a longer incubation time, the tryptic activity affects the cells and therefore the temperature should be minimized to reduce the effect triptych. The easy way to reduce damage to the cells during exposure is to soak the tissue with trypsin at 4 ° C for 6-18 hours to allow penetration of the enzyme with little triptych activity followed by exposure of 20 to 30 minutes at 37 ° C for disintegration.

Rats (approximately 200 g) were anesthetized by inhalation of chloroform, and the liver was isolated by a horizontal incision on the ventral surface near the diaphragm of the animal.

The liver of dead time does not last more than 2 to 3 min. The liver was kept in saline phosphate buffer solution (PBS) at 4 ° C until use. The experimental environment was sterilized.

0.375 g liver was removed and then cut with the surgical blade into small pieces; these pieces were washed with PBS and then transferred to other Petri dishes and unwanted tissue such as fat tissue and necrotic tissue were removed. Liver pieces were washed once more with PBS; transferred to a falcon tube in which they are soaked in 0.25% crude trypsin in RPMI for 18 hours at 4 ° C. Trypsin is added following the ratio of 10 ml per 1 g of tissue.

After incubation, the liver pieces were exposed at 37 ° C for 30 minutes in 1 ml of warm RPMI per 100mg of tissue. In order to completely disperse the tissue, a gentle pipetting is performed, then the solution is filtered through a sterile muslin cloth to remove the larger pieces. Cells were preserved for experiments and stored in common crop in 5% dimethyl sulfoxide

(DMSO) used as cryoprotectant, added to the cell to reduce the formation of ice and thereby prevent cell death in during the freezing process, and the cells were frozen at -80 °

C.



Figure 16: liver cells of rats Photo isolated and stored in culture in 5% DMSO (Photo Amos Oluwagbemiga S., 2015).

Extraction and thin layer chromatography of cellular lipids Lipid extraction

The lipid extraction method is based on the method called solvent gradient used by Stephan et al. (2004) for extracting mycobacterial lipids. It has been adapted for hepatocytes. Four successive extractions are carried out on the mêméchantillon cells. The extraction solvent is a methanol / chloroform increasingly enriched in chloroform [11; 29].

- Cellular lipids were extracted by four successive extractions of a day, stirring at 250 rpm. The lipid extract is recovered after separation of the cells; adding an aqueous solution of NaCl 0.9% (volume proportional to CHCl₃) afforded the two phases [30; 31] and the cells are then ready to undergo the next extraction.

- The proportion of solvents chloroform (CHCl₃) and methanol (CH₃OH) are different depending on the extracting step:

o First extraction: CHCl₃ / CH₃OH (1V: 2V)

o Second extraction: CHCl₃ / CH₃OH (1V: 1V)

o Third extraction: CHCl₃ / CH₃OH (2V: 1V)

o Fourth extraction: CHCl₃ / CH₃OH (2V: 1V)

A volume V = 15 ml for 500 mg of cells

- All the organic phases were mixed and the solvent is evaporated in a Rotavapor and under hood.

- The lipid extract is resuspended in a volume of chloroform (1 ml solvent per 1 g of lipids).

- Part of the lipid extract is used for analysis and the other is stored at -80 ° C.

The sample was stored at - 80 ° C is evaporated under a stream of nitrogen and then resuspended in chloroform to know the precise volume before use analysis of lipids [11].



Figure 17: Phase separation photos (below) containing chloroform lipids by settling.



Figure 18: Photograph of the lipids extracted from solutions in solvent (chloroform) evaporates.

Thin layer chromatography (TLC) of cellular lipids This is the separation of different classes of silica gel lipids by thin layer chromatography. CCM is based on electrostatic interaction / hydrogen bonding. The principle of "like attracts like that," often encountered in chemistry still helps explain here the nature of the phenomena involved. It is based on the differential adsorption phenomena: a mobile phase consisting of solvent progresses along a stationary phase (silica) fixed on a plate. TLC takes place in three stages: preparation of the tank, preparation of the plate, and elution. The different cellular lipids are divided into two main groups: phospholipids and neutral lipids. They are separated on two different plates. [11]

Separation of neutral lipids by TLC

As their name suggests, the neutral lipids are not charged compounds. They include: the free or esterified fatty acids, mono-, di- and triglycerides and free and esterified sterols. They are separated using low- polar solvents: hexane, ethyl ether, ... The separation of neutral lipids is done first by a migration in a hexane solvent: ethyl ether: glacial acetic acid (40: 10: 1 v / v / v) up to half laplaque. The plate is dried and then is

performed a second migration up to 1 cm from the upper edge of the plate with hexane alone [11; 31].

- Preparation of the tub (mobile phases)

- Eluent E1: it corresponds to the mixture M1, the following solvents: hexane: ether: ethyl: glacial acetic acid(40: 10: 1 v / v / v) [11; 31].

- Eluent E2: it is the only hexane. [11]

The two eluents are placed in two different tanks C1 and C2 at a height of about 0.5 cm from the bottom of the vessel; and the tanks are closed so that they are saturated with vapor eluant for 10 to 15 min.

- Preparation of the stationary phase

Silica plate (Silica gel 60, Merck, France) of dimensions 10 × 10 cm, is drawn in pencil, a line about 1 cm from the bottom edge and parallel thereto; then placed on the line, using capillary tubes and 0.5 to 1 cm apart, 10 mL of each lipid sample (control, sample 1, sample 2 ...) [11; 31].

- Elution or development of the chromatogram

- The plate is placed slightly oblique position in the C1 tank is then closed. The solvent which covers the bottom rises along the plate by capillary action.

- Migration is stopped when the solvent front reaches about half of the plate.

- The plate is then placed in the second tank C2 after drying and migration is stopped when the solvent frontreaches about 1cm of the upper end of the plate.

- The level reached by the solvent (front line) is marked by a thin line and the plate is dried in the open air orusing a drier [11; 31].



Figure 19: Photographs showing a first eluting in hexane: ethyl ether: glacial acetic acid (40: 10: 1 v / v / v) up to half of the plate (left) and a second elution in hexane until a cm from the upper edge of the plate (right).

2.3.4.2.1. Separation of phospholipids by TLC

Phospholipids are polar molecules separated using highly polar solvent combinations: water, methanol, chloroform ... The migration was carried out at once with a mixture: Chloroform / methanol / ammonia (32.5: 12.5: 2). Such proportions are at the limit of the two-phase mixture. We must remain in monophasic [11; 31].

- Preparation of the vessel (mobile phase)

Pharma Research [2021]13(1)1-4

The eluent is the mixture of the following solvents: chloroform / methanol / ammonia (32.5: 12.5: 2). The eluent was placed in a C3 tank at a height of about 0.5 cm from the bottom of the vessel; then closes the tank so that it is saturated with steam eluent during 10 to 15 min [11; 31].

• Preparation of the stationary phase

Silica plate (Silica gel 60, Merck, France) of dimensions 10 × 10 cm, is drawn in pencil, a line about 1 cm from the bottom edge and parallel thereto; then placed on the line, using a capillary tube and 0.5 to 1 cm apart, 10 mL of each lipid

sample (control, sample 1, sample 2, ...)

• Elution or development of the chromatogram

- The plate is placed slightly oblique position in the C3 tank is then closed. The solvent which covers the bottom rises along the plate by capillary action.

- Migration is stopped when the solvent front reached approximately 1 cm in the upper end of the plate.

- The level reached by the solvent (front line) is marked by a thin line and the plate is dried in the open air or using a drier [11; 31].



Figure 20: Photograph showing elution in chloroform / methanol / ammonia (32.5: 12.5: 2) to one cm from the top of the plate.

Revelation of lipids on TLC plates

The detection is carried out by spraying a copper sulfate solution II 10% (10 g CuSO₄ 5H₂O in 100 mL of H₃PO₄ 8% (v / v)). Once the dry plate, it is placed for about 15 min in an oven at 150 ° C min. The presence of spots indicates the presence of lipids in question. [11]

II. RESULTATS

The extraction of Return is calculated using the formula: $R = \text{Mass of l'extract} / \text{Mass of powder of bark} \times 100$

Extract	Yield,%	Color	Appearance
Decoction / water	R= 27,3×100/300 R= 9,1	Red wine	Powder

The yield of 9.1% obtained for extraction is low

Mass lipids extracted

Essay	Témoi n (0mg/ Kg)	Lot 1 (50mg /K g)	Lot 2 (100mg / Kg)	Lot 3 (200mg / Kg)	Lot 4 (300mg / Kg)	Lot 5 (400mg / Kg)
Avera geMas s of lipid extrac ts	0.132 mg	0.196 mg	0.237 mg	0.131 mg	0.084 mg	0.152 mg

Table showing the average masses of lipids extracted from Rat Hepatocytes

3.3. Action of the aqueous extract of the bark of *S.birrea* Profile neutral lipids rat hepatocytes



II- T 50 100 200 300 400

Figure 21: Photograph of the plate to UV 366 nm before revelation with CuSO_4 II, showing the migration of different neutral lipids. (Photo of Oluwagbemiga Amos S., 2015)



T 50 100 200 300 400

Figure 22: Picture of the plate with UV 254 nm before revelation with CuSO_4 II, showing the migration of different neutral lipids. (Photo of Oluwagbemiga Amos S., 2015)



T 50 100 200 300 400

Figure 23: Photo of the plate after development with CuSO_4 II, showing the migration of different neutral lipids. (Photo of Oluwagbemiga Amos S., 2015)

Rats batches treated with the extract, it was found that the daily administration of different doses of ethyl extract from the bark of *Sclerocarya birrea* relative to the control group during 14 days influenced qualitatively and quantitatively different types of lipids neutral in cells. The figure 23; 24 and 25 show the pictures of the plate obtained UV 260 nm respectively, UV 340 nm and after revelation with CuSO₄ II after a first migration with hexane / diethyl ether / glacial acetic acid (40: 10: 1 v / v / v) up to half of the plate, then a second migration with hexane alone to 1 cm from the edge of the plate. This allows to separate the different neutral lipids. The first deposit corresponds to the negative control and the other are respectively the doses of 50mg / kg, 100mg / kg, 200mg / kg, 300mg / kg and 400mg / Kg of ethyl extract from the bark of *S. birrea*. Whereas in the neutral lipids in their entirety is found that:

- To witness so there's about five spots five neutral lipids

- For doses of 50mg / kg and 100mg / kg was also about five spots so five neutral lipids
- For doses of 200mg / kg, 300mg / kg and 400mg / kg was about seven spots so seven neutral lipids

Considering each lipid, note that:

- The spots at the witness are very low, so each lipid in question is small amount compared to other higher doses.

- The spots at doses of 50mg / kg and 100mg / kg are as low as those of the control relative to the other, so the amount of each of the lipids in these doses is almost the same as in the control over witness

- The spots at doses of 200mg / kg, 300mg / kg and 400mg / kg are sharper, darker compared to the control and the first doses so the amount of each lipid at these doses is greater by compared to the control and all other doses.

In addition, we note that at each deposit there are more heavy lipids (in low spots) as light lipids (on high spots).

III. ACTION ETHYL EXTRACT OF THE BARK OF SCLEROCARYA BIRREA PROFILE PHOSPHOLIPIDS RAT HEPATOCYTES




Figure 24: Picture of the picture plate of the plate to UV 366 nm before revelation with CuSO₄ II, showing the migration of various phospholipids. (Photo of Oluwagbemiga Amos S., 2015)



Figure 25: Photograph of the picture plate of the plate with UV 254 nm before revelation with CuSO₄ II, showing the migration of various phospholipids. (Photo of Oluwagbemiga Amos S., 2015)




Figure 26: Photo of the plate after development with CuSO_4 II, showing the migration of different phospholipids (Photo of Oluwagbemiga Amos S., 2015).

Daily administration of different doses of the aqueous extract of the bark of *Sclerocarya birrea* rats batches treated with the extract compared to the control group for 14 days influenced phospholipids in cells. Figures 26; 27 and 28 show the pictures of the plate obtained UV 260 nm respectively, UV 340 nm and after revelation with CuSO_4 II after migration in a mixture: Chloroform / methanol / ammonia (32.5: 12.5: 2) until to 1 cm from the edge of the plate. Indeed this migration has effectively separate the different phospholipids. The first deposit corresponds to the negative control and the other are respectively the doses of 50mg / kg, 100mg / kg, 200mg / kg, 300mg / kg and 400mg / kg of aqueous bark extract *Sclerocarya birrea*. Considering the phospholipids in their entirety shows that:

- For the control there are about five spots so five phospholipids.
 - For doses of 50mg / kg, 100mg / kg, 200mg / kg, 300mg / kg and 400mg / kg was also about five spots so five phospholipids.
- However considering each lipid, note that:
- The spots at the witness and doses of 50mg / kg and 100mg / kg are very small, so each of the phospholipids in question is small amount compared to other higher doses.
 - The spots at doses of 200mg / kg, 300mg / kg and 400mg / kg are sharper, darker compared to

the control and to the first two doses so the amount of each of the phospholipids at these doses is greater by compared to the control and all other lower doses.

Furthermore, we note that at each submission it has more heavy fat y (down spots) only minor lipids (high in sp

IV. D IS C U S SI O N

The extraction yield is 9.1%. This result shows that the dry peel *Sclerocarya trunk birrea* contain active ingredients which are soluble in water, therefore extractable by the latter. In addition, the yield is low compared to that of Attakpa et al (16%) [32] and Ivonne Gisèle Makom Ndifossap et al (16%) [33], who conducted the aqueous extraction by maceration. These differences could be explained either by the extraction methods or by either solvents.

The administration of aqueous extract of the bark of *Sclerocarya birrea* at various doses: 50-400mg / kg shows on the photos 23, 24, 25

Pharma Research [2021]13(1)1-4

there is an increase in neutral lipids in cells having undergone the action of *S. birrea* large quantities: 200-400mg / kg than in the cells of the negative control and rats having undergone the action at low doses (50-100mg / kg); and this increase is proportional to the dose of *S. birrea* administered. So

we can deduce that *S. birrea* stimulates either the neutral lipid biosynthesis process or be neutral lipids are from degradation of membrane phospholipids.

- Each neutral lipid biosynthesis of processes emphasize the stimulation of several voice signaling, the synthesis of several enzymes and therefore the transcription of several genes. The idea that the *S. birrea* molecule alone can trigger all these processes seems unlikely. But these processes could be triggered by the presence of lesions in the membrane to compensate for losses of phospholipids.

- In the phospholipids biosynthesis process, neutral lipid (mono-, di- and tri- glycerides) serve as precursors [8; 34]. So conversely, in case of degradation of the phospholipids could have release neutral lipids.

Thus, the increase of neutral lipids through the action of *S. birrea* come from degradation of membrane phospholipids.

In addition, administration of the aqueous extract of the bark of *S. birrea* the 50 and 100mg / kg doses shown in photos 23, 24 and 25 there is no significant change in the neutral lipids profile compared the negative control. So in doses 50 and 100 mg / Kg there can be no degradation of phospholipids.

Photos 26, 27 and 28 show the action of the aqueous extract of the bark of *Sclerocarya birrea* at various doses: 50-400mg / kg of phospholipids. We see that there an increase in phospholipids in doses of 200,300 and 400 mg / kg compared to the negative control, while in doses 50 and 100 mg / kg there was no significant change compared to the negative control. Phospholipids are specifically membrane molecules, their strong presence in the lysates under the action of *Sclerocarya birrea* compared to the negative control may be explained by the fact that even before cell lysis by extraction solvents extract *Sclerocarya birrea* had induced the lysis of the plasma membrane; and this lysis is nothing but the result of necrosis. This result allows us to understand that the degradation of membrane lipids cited above is actually the result of necrosis.

CONCLUSION

We investigated the effects of this plant's bark aqueous extract on the rat liver's lipid profile. *Sclerocarya birrea* seems to be hazardous at high dosages (200 mg/kg, 300 mg/kg, and 400 mg/kg) according to the results of this investigation. The fact that the lipid profile did not alter considerably at the 50 mg/kg and 100 mg/kg doses relative to the negative control suggests that our study may help

us avoid this toxicity. *Sclerocarya birrea* aqueous extract is hazardous at concentrations of 200 mg/kg or above.

With these findings, traditional healers will have more information to help their patients avoid liver damage while using *Sclerocarya birrea*, which bodes well for the future. We think our findings are noteworthy because they pave the way for new avenues of investigation in the field of experimental biology.

REFERENCES

- [1]INTEVHELENI THOVHOGI (2009) examined the effects of sclerocaryabirrea stem bark extracts on the activity of certain carbohydrate metabolising enzymes associated with diabetes.
- [2]. In 2011, GOUWAKINNOU et al. published an article in the Journal of Ethnobiology and Ethnomedicine. In 2010, Bintou Maïga studied the phytochemical composition, anti-radical activity, and subchronic toxicity of the leaves of *Sclerocarya birrea* (A. Rich) Hoscht (Anacardiaceae), which are used in traditional diabetes treatment in Mali. The thesis was completed at the University of Bamako and was published in the first part of the thesis, pages 44-50.
- [3]. English Wikipedia
- “Contribution à l'étude des produits de cueillette (PFC): Potentiel et utilisations dans le terroir d'Ambandé (région de Kaolack) au Sénégal” [4] by Mamadou Diane. Dakar's Cheikh Anta Diop University "Phytosociological and chorological approaches to phytogeography: a meso-scale study in Benin" (2006) by Adomou AC, Sinsin B, and Van der Maesen LJG. (Plant Systematics and Geography, 76:155-178.)
- [5]. "Cellular membranes" by B. Lahlou et al.8. The Foundation (2010). The case of disseminated neoplasie in bivalves: an investigation of the relationships between lipid membrane compositions and cellular functions. Finalised May 14, 2010. University of the Western Hebrides. Pages 40–69.
- [6]. Thou, M. H. (2010). Under stress, changes occur in the membrane lipid content of lactobacilli. Exploration of the physiological function of cyclic fatty acids in two models: *Oenococcus oeni* ATCC-BAA1163 and *Lactococcus lactis* MG1363. Pages 5–30 of the thesis submitted to the Institut Universitaire de La Vigne et du Vin at the Université de Bourgogne.
- [7]. 2010 by DELPHINE CASTAGNE. Research on the interactions between cyclodextrines and biological or liposomal membranes. Documentation from the University of Liège. 141P.
- [8]. "Marlene Cot" from 2006. Physiological studies of the adaptability and resistance of the yeast *Saccharomyces cerevisiae* in the context of extensive ethanol production. Page number: 71–87 from the Toulouse National Institute of Applied Sciences' thesis. The authors of the article are Bereziat, Chammaz, Cordard, and Wolf (1988). The many roles of cell membrane phospholipids. *Medecine/Sciences* 1988; 4: 8–15 Cnrs, UA 524–, Laboratoire de Biochimie, CHU St-Antoine, 27, rue Chaligny, 7501 2 Paris, France. 13. MATTHIEU S. founded Cours-Pharmacie and serves as its chief editor [9]. 15 from the Lessire M1 MB 2007 PDF file. VEE, LE (2006). Improving in vitro study models

Pharma Research [2021]13(1)1-4

and investigating mechanisms that regulate their expression: proteins involved in membrane transport of xenobiotics. 18P. 16 GRONDIN, M. (2008). Winter wheat proteins: novel cryoprotective agents for rat liver cells. Chapter II of the thesis from the University of Quebec, pages 8–18. 17. NATAF S. Histology: le-foie-et-les-voies-biliaires.htm. The thesis was written by S. Lyon Civil Hospitals and the University of Lyon 9P.

AHOKPE, M. (2012), [10]. Endocrine course 57P. "AHOKPE M. (2012)" [11]. The digestive system, the liver, and the bile ducts are covered in histology's Organs, Systems, and Appearances chapter 1. Level PCEM2-DCEM157P at the Pierre and Marie Curie Medical School. Section 6 of the NATAF S. Course in Medicine—Anatomy, PCEM1, Ostéopathie—Le foie [12].12P. in the year 2010 (SARF, F. S.). Simulation of the transport process of a class of substances across cellular membranes using biochromatographic and thermodynamic methods. Page 35–37 of the thesis from the University of Franche-Comté. In 2007, the authors Catala, André, Katsanis, and Poirier published a paper. Human anatomy: systems, organs, and tissues. Université Pierre et Marie Curie, Service d'Histologie et Embryologie; Cours de PCEM2 - DCEM1. 11P. As of November 2005, the source is KOUKOU A.S.I. OMEGINE. Investigating the hepatocellular calcium signals induced by various bile acid, prostaglandin, and glucagon agonists that control bile secretion and hepatic glucose synthesis. Thesis presented at the XI ORSAY University of Paris; pages 1-16. the work of M. C. Takin (2014). Pharmacodynamics of the bioactivity of ethyl Khayasenegalensis extract on liver and kidney functions in the rabbit (*Orytholagus cuniculus*). Chapter I, pages 7–17, Thesis from the University of Abomey-Calavi, 01 BP526 Cotonou, Benin. [17]. Published by Prajapati and Patel in 2011. Cold trypsinization for rat hepatocyte separation and hot sds/phenol extraction for total RNA isolation...Publication date: 2011 Jan 3; volume: 3, issue: 1, pages 3–3. cited as [18]. This was published in 1989 by Sambrook, Fritsch, and Manitis. Genomic Recombination. Chapter III: Experimental Methods in a Laboratory Manual Published by Cold Spring Harbour Laboratory Press in New York. [19]. The methods of nucleic acid separation and analysis described by MEDRAOUI L. in Adn, Arn, and Plasmides. Section 1: Molecular Biology, Module M21, Element 2. Research Centre for Molecular Biology and Microbiology at the Rabat Faculty of Science and the University of Mohamed V. Agdal [20]. In 2008, Fernando Streit wrote. The physiological state and cryotolerance of *Lactobacillus delbrueckii* subsp. *bulgaricus* cfl1 are affected by harvest and concentration conditions. Agro Paris Tech theses from the Institut des Sciences et Industries du Vivant et de l'Environnement (pages. 91–95). Clément, Segu-real, Savage, Kleinman, and Yamada (1990) were the authors of the cited work. Multiple receptors mediate laminin's attachment to hepatocytes. Developmental biology and abnormalities laboratory. Bethesda, Maryland is home to the National Institute of Health's National Institute of Dental Research. 8092. 8P.

The number 22. Author: Sonoia do Carmo (2009). Investigation of human apolipoprotein E in response to stress. Quebec University thesis, Montréal, Quebec, Canada, 33–38 pages. The number 23. This sentence was written by Kolawoole, Olaniran, Ayankunle, and Kolawoole in 2012. Choyasenegalensis Stem Bark Aqueous Extract's Anti-hyperglycemic Effect in Wistar Rats. Citation: European Journal of Medicinal Plants; 2(1): 66-73. [24]. (2014) by Montemo G.F., Tehoua L., Ouattara H., and Yapi A. A study comparing the hypoglycemic effects of *Sclerocaryabirrea*, *Khayasenegalensis*, *Heliotropiumindicum*, and *Ocimum gratissimum* on rats. Published in the American Journal of Bioscience, volume 2, issue 2, pages 60–63. [25]. A study conducted by ATTAKPAET AL. (2015) found that a stem-bark extract from *Sclerocarya Birrea* (Anacardiaceae) stimulates the liver's protein kinase Akt and AMPK pathways in a diet-induced obesity mouse model. [26]. (2012) by IVONNE GISÈLE MAKOM NDI FOSSAP and colleagues. In diabetic rats, stem-bark extract from *Sclerocaryabirrea* (Anacardiaceae) improves glucose tolerance and insulin production in response to glucose stimulation [27]. The work of J. B. M. Ratray, included in *Microbial Lipids*, edited by C. Ratledge and S. G. Wilkinson (1988), spans pages 555-698. Jovanovich & Harcourt